


# Non-Hermitian ensembles and Painlevé critical asymptotics



Alfredo Deaño & Nick Simm

University of Kent, University of Sussex

A. Deano-Cabrera@kent.ac.uk, N. J. Simm@sussex.ac.uk



## Normal matrix model

We are interested in the normal random matrix model defined by

$$dP_N(z_1, z_2, \dots, z_N; t) = \frac{1}{Z_N(t)} \prod_{j=1}^N (z_j)^2 e^{-NV_t^{(s)}(z_j)} dA(z_j);$$

with  $z_j \in \mathbb{C}$  and potential

$$V_t^{(s)}(z) = jz^{2s} - t(z^s + \bar{z}^s); \quad s \in \mathbb{N};$$

The eigenvalues  $z_1, \dots, z_N$  display an interesting behaviour:

Figure 1: The limiting eigenvalue distribution is supported on the interior of the orange curves. Here  $s = 11$  and  $t = t_c - 0.1$  (left),  $t = t_c$  (centre) and  $t = t_c + 0.1$  (right). At the special value  $t = t_c$ , the support becomes disconnected.

In this poster our goal is to investigate the partition function  $Z_N(t)$  near the critical value  $t = t_c = 1 = \frac{1}{2s}$ .

## Reduction to the Ginibre ensemble

The first observation is that we can use symmetry to write  $Z_N(t)$  as an average over the Ginibre ensemble:

$$Z_{Ns}(t) = c_{N;s} \int_{\mathbb{C}^N} \prod_{j=1}^N Z_N^{(j)}(x); \quad j := 2, 1, \frac{l+1}{s};$$

where

$$Z_N^{(j)}(x) = \int_{\mathbb{C}^N} \prod_{j=1}^N (z_j)^2 e^{-Njz_j^2} e^{-Njz_j^2} dx_j e^{-Njz_j^2} dA(z_j); \quad x := t^{\frac{1}{2s}};$$

Criticality now corresponds to the spectral variable  $x$  colliding with the boundary of the circular law (i.e.  $|x| = 1$ ). When  $|x| < 1$  (sub-critical), the asymptotics were obtained in [2].

## Painlevé and non-Hermitian matrix integrals

Our main result for finite  $N$  characterizes the partition function as a solution of the  $\sigma$ -form of Painlevé V.

**Theorem 1.** The 'reduced' partition functions  $Z_N^{(j)}(x)$  can be written as

1. An average over the CUE:

$$Z_N^{(j)}(x) = c_N \int_{\text{CUE}} \prod_{j=1}^N e^{-\frac{i}{4}j|1 + e^{i\frac{j}{2}}|} e^{Nx^2 e^{ij}};$$

2. The  $\sigma$ -form of Painlevé V:

$$Z_N^{(j)}(x) = c_N \exp \int_0^{Nx^2} \frac{y_N(t) + \frac{N}{2}}{t} dt$$

where  $y_N(t)$  satisfies the equation

$$(t^{\theta})^2 \left[ t^{\theta} + 2(t^{\theta})^2 + (N - \theta)^2 \right]^2 + 4(t^{\theta})^2 (t^{\theta} + N) = 0; \quad (1)$$

with initial condition

$$y_N(t) = \frac{N}{2} + \frac{t}{2}; \quad t \rightarrow 0;$$

The first part above can be arrived at by a judicious inspection of formulas in [2]. Then the second part is a consequence of the first due to results of Forrester and Witte '02.

## Large $N$ asymptotic results

Asymptotic results for related orthogonal polynomials have been studied in various works, but the critical case only very recently in [1]. For the partition function, we obtain:

**Theorem 2.** If  $k = 2k$ , where  $k \in \mathbb{N}$ , then for

$$|x| = 1 - \frac{u}{N}; \quad u \in \mathbb{R}$$

we have the following asymptotics:

$$\frac{Z_N^{(2k)}(x)}{E_{N;k}} = \exp \left( \rho \int_{\mathbb{C}} \frac{1}{Nku} \int_{\mathbb{C}} v(z) d\mu(z) \right) (1 + o(1)); \quad N \rightarrow \infty;$$

uniformly for  $u$  in compact subsets of  $\mathbb{R}$ , where  $E_{N;k}$  is a completely explicit pre-factor. The function  $v$  satisfies the  $\sigma$ -form of the Painlevé IV equation:

$$(v^{\theta})^2 + 4(v^{\theta})^2(v^{\theta} + k) - (sv^{\theta} - v^{\theta})^2 = 0; \quad (2)$$

subject to the boundary condition

$$v(s) = ks - \frac{k}{s} + O(s^{-3}); \quad s \rightarrow \infty;$$

We believe this result persists to non-integer  $k$ , indeed a naive rescaling of equation (1) reproduces exactly the Painlevé IV in (2). The advantage of integer  $k$  is the duality (Forrester and Rains '08):

$$\frac{Z_N^{(2k)}(x)}{E_{N;k}} = jx^{2Nk+2k^2} \prod_{j=1}^N e^{-\frac{1}{N}jx^{2j}} \left( 1 + \frac{r_j}{N} \right)^N \int_{\mathbb{C}} (F)^2 dF; \quad (3)$$

making  $N \rightarrow \infty$  asymptotics easy to compute. For  $k$  not integer, we use Riemann–Hilbert techniques.